

# HPCS CUSTOM STOCKER R

## MEDICATED

### For Growing Beef Cattle

For increased rate of weight gain in growing cattle on pasture or dry lot (stocker, feeder; dairy and beef replacement heifers).

#### ACTIVE DRUG INGREDIENT

Monensin (as Monensin Sodium) . . . . . 1,200 g/ton

#### GUARANTEED ANALYSIS

Calcium (Ca), minimum . . . . .	15.30%
Calcium (Ca), maximum . . . . .	18.30%
Phosphorus (P), minimum . . . . .	3.50%
Salt (NaCl), minimum . . . . .	18.00%
Salt (NaCl), maximum . . . . .	21.60%
Magnesium (Mg), minimum . . . . .	1.50%
Potassium (K), minimum . . . . .	0.30%
Copper (Cu), minimum . . . . . ppm	2,100
Selenium (Se), minimum . . . . . ppm	26.50
Zinc (Zn), minimum . . . . . ppm	7,600
Vitamin A, minimum . . . . . IU per lb	270,000
Vitamin D <sub>3</sub> , minimum . . . . . IU per lb	27,000
Vitamin E, minimum . . . . . IU per lb	270

#### FEED INGREDIENTS

Calcium Carbonate, Salt, Monocalcium Phosphate, Dicalcium Phosphate, Corn Distillers Dried Grains with Solubles, Molasses Products, Magnesium Oxide, Zinc Amino Acid Complex, Manganese Sulfate, Zinc Oxide, Copper Amino Acid Complex, Basic Copper Chloride, Sodium Selenite, Potassium Chloride, Vitamin E Supplement, Vitamin A Acetate, Ethylenediamine Dihydroiodide, Vitamin D<sub>3</sub> Supplement, Cobalt Carbonate, Vegetable Fat (preserved with BHT)

#### FEEDING DIRECTIONS

Thoroughly mix HPCS Custom Stocker R Medicated with grain or roughage before feeding to growing beef cattle. Feed 4 ounces of HPCS Custom Stocker R Medicated mixed in 1 lb or more of feed on 90% dry-matter basis to supply 150 mg of Monensin per head per day. During the first 5 days, mix 2.67 ounces of HPCS Custom Stocker R Medicated in at least 1 lb or more of feed so cattle receive no more than 100 mg of Monensin per head per day.

Do not exceed 4 oz per head per day so that the intake of selenium does not exceed the legal maximum of 3 mg per head per day.

CAUTION: Follow label directions. The addition to feed of higher levels of this premix containing selenium is not permitted.

WARNING: CONTAINS HIGH LEVELS OF COPPER. DO NOT FEED TO SHEEP.

CAUTION: Do not allow horses or other equines access to feed containing Monensin. Ingestion of Monensin by horses has been fatal. Monensin medicated cattle and goat feeds are safe for use in cattle and goats only. Consumption by unapproved species may result in toxic reactions.

Do not feed undiluted. Feeding undiluted or mixing errors resulting in high concentrations of Monensin has been fatal to cattle and could be fatal to goats. Do not exceed the levels of Monensin recommended in the feeding directions, as reduced average daily gains may result. If feed refusals containing Monensin are fed to other groups of cattle, the concentration of Monensin in the refusals and the amount of refusals fed should be taken into consideration to prevent Monensin overdosing.

This product was made in a feed facility that does not manufacture products containing animal proteins prohibited in ruminant feed.

This feed was produced in a facility participating in the American Feed Industry Association's Safe Feed/Safe Food Certification Program. Go to [www.afia.org](http://www.afia.org) for details.



Manufactured By:  
Provimi North America, Inc.  
Brookville, OH 45309



Rev. 10-25-17

F111135055

**Net Weight 50 lb (22.68 kg)**

Red

